

HOSTED BY




ELSEVIER

Contents lists available at ScienceDirect

China University of Geosciences (Beijing)

Geoscience Frontiers

journal homepage: www.elsevier.com/locate/gsf

Research paper

The giant Jiaodong gold province: The key to a unified model for orogenic gold deposits?

David I. Groves^{a,b}, M. Santosh^{b,c,d,*}^a Centre for Exploration Targeting, UWA, Crawley 6009, WA, Australia^b Centre for Tectonics Resources and Exploration, Department of Earth Sciences, University of Adelaide, SA 5005, Australia^c State Key Laboratory of Geological Processes and Mineral Resources, China University of Geosciences, Beijing 100083, China^d Division of Interdisciplinary Science, Faculty of Science, Kochi University, Kochi 780-8520, Japan

ARTICLE INFO

Article history:

Received 2 July 2015

Received in revised form

27 August 2015

Accepted 30 August 2015

Available online 14 September 2015

Keywords:

Jiaodong province

Orogenic gold

Metamorphic fluids

Subduction zones

Tectonics

ABSTRACT

Although the term orogenic gold deposit has been widely accepted for all gold-only lode-gold deposits, with the exception of Carlin-type deposits and rare intrusion-related gold systems, there has been continuing debate on their genesis. Early syngenetic models and hydrothermal models dominated by meteoric fluids are now clearly unacceptable. Magmatic-hydrothermal models fail to explain the genesis of orogenic gold deposits because of the lack of consistent spatially – associated granitic intrusions and inconsistent temporal relationships. The most plausible, and widely accepted, models involve metamorphic fluids, but the source of these fluids is hotly debated. Sources within deeper segments of the supracrustal successions hosting the deposits, the underlying continental crust, and subducted oceanic lithosphere and its overlying sediment wedge all have their proponents. The orogenic gold deposits of the giant Jiaodong gold province of China, in the delaminated North China Craton, contain ca. 120 Ma gold deposits in Precambrian crust that was metamorphosed over 2000 million years prior to gold mineralization. The only realistic source of fluid and gold is a subducted oceanic slab with its overlying sulfide-rich sedimentary package, or the associated mantle wedge. This could be viewed as an exception to a general metamorphic model where orogenic gold has been derived during greenschist- to amphibolite-facies metamorphism of supracrustal rocks: basaltic rocks in the Precambrian and sedimentary rocks in the Phanerozoic. Alternatively, if a holistic view is taken, Jiaodong can be considered the key orogenic gold province for a unified model in which gold is derived from late-orogenic metamorphic devolatilization of stalled subduction slabs and oceanic sediments throughout Earth history. The latter model satisfies all geological, geochronological, isotopic and geochemical constraints but the precise mechanisms of auriferous fluid release, like many other subduction-related processes, are model-driven and remain uncertain.

© 2015, China University of Geosciences (Beijing) and Peking University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Groves et al. (1998), following Gebre-Mariam et al. (1995), defined the term orogenic gold deposit to obviate the necessity to refer to a wide variety of terms for a coherent group of commonly vertically-extensive, gold-only deposits that formed in broad thermal equilibrium with their wallrocks from low-salinity H₂O-CO₂ ore fluids at depths from 2 to 15 km, and arguably 20 km in the

crust (Groves, 1993; Kolb et al., 2015). This term has been widely accepted (e.g., Goldfarb et al., 2001, 2005, 2014; Bierlein et al., 2006), although there is still some discussion on terminology (e.g., Phillips and Powell, 2015), and a heated debate on the genesis of orogenic gold deposits is ongoing. Goldfarb and Groves (2015) provided an exhaustive review of these genetic models and the various geological, geochemical, isotopic and fluid-inclusion constraints on these models. This review is used, comprehensively, to briefly summarize these models with a view to provide a holistic, coherent and unified model for orogenic gold deposits of all ages, in a similar way to development of coherent minerals-system models for other mineral deposit groups. The deposits of the giant Jiaodong

* Corresponding author.

E-mail addresses: msantosh.gr@gmail.com, santosh@cugb.edu.cn (M. Santosh).

Peer-review under responsibility of China University of Geosciences (Beijing).

orogenic gold provinces are emphasized as the key to development of the all-embracing model for ore fluid and metal source.

It is recognized that (reduced) intrusion-related gold systems or (R)IRGDs (e.g., Thompson et al., 1999; Lang et al., 2000; Baker, 2002) formed from a similar ore fluid to the orogenic gold deposits, but that they differ in that the ore systems are zoned around causative intrusions due to thermal disequilibrium with the wall rocks (e.g., Hart et al., 2002). They are, however, a rare group of largely un-economic deposits (e.g., Goldfarb et al., 2005; Goldfarb and Groves, 2015), and are not discussed further here. Furthermore, although the Carlin gold deposits also formed from low-salinity H₂O–CO₂ fluids (e.g., Cline et al., 2005), they are quite distinctive from orogenic gold deposits in a number of features (Goldfarb and Groves, 2015; Groves et al., 2016), and are not discussed below.

2. Potential fluid and metal sources for orogenic gold deposits

Kerrick (1983) was arguably the first to assess the various models for what are now termed orogenic gold deposits, listing syngenetic-exhalative, magmatic-hydrothermal (tonalite-, lamprophyre- or oxidised magma-associated), and metamorphic (regional metamorphic devolatilization, lateral secretion, mantle/granulitization) models as the major suggested genetic concepts for fluid and metal generation. A meteoric fluid model was added by Nesbitt (1991). All of these models are shown schematically in Fig. 1.




Figure 1. Schematic representation of the variety of previously proposed (mostly non-viable) models for gold and fluid sources in the crust: from meteoric water circulation and lateral secretion, magmatic-hydrothermal fluid exsolution from various granite types, to granulitization and metamorphic devolatilization processes. Syngenetic-exhalative model is not shown, but could be represented by the hot springs at surface in the figure. Figures from Groves et al. (1998) and Goldfarb et al. (2005) used as a base for this figure.

Goldfarb and Groves (2015) discussed each of these models in detail with exhaustive references to individual examples in places. The following brief discussion of the less-viable models is adapted from Goldfarb and Groves (2015), and then followed by a more thorough discussion of the more-viable, more generally-accepted models.

Early syngenetic-exhalative models (e.g., Hutchinson and Burlington, 1984; Hutchinson, 1987) were shown to be inconsistent with field evidence that demonstrated the deposits were structurally-controlled, syn- to late-metamorphic deposits with stratiform BIF-hosted deposits formed by sulfidation of magnetite (e.g., Phillips et al., 1984). Similarly, meteoric fluid models have been shown to be based on invalid calculations and interpretations of stable isotope data largely derived from fluid inclusions, as summarized by Goldfarb and Groves (2015).

Various magmatic-hydrothermal models were in vogue for a variety of mineral deposits from about 1900 to 1960, and have been proposed for orogenic gold deposits over the past 40 years or so by a number of authors, most recently including Mueller (1992), Walshe et al. (2003), Wall et al. (2004), Hall and Wall (2007), Neumayr et al. (2007), Bath et al. (2013) and Helt et al. (2014). Goldfarb and Groves (2015, and references therein) discussed these models at length for a number of specific examples and rejected the magmatic-hydrothermal concept as a viable unifying model for orogenic gold deposits. In general, granitic intrusions may be pre-, syn- or post-gold in the same terranes (e.g., Hughes et al., 1997; Goldfarb et al., 2008), or even absent in some, for example in the Otago gold province of New Zealand. In most cases where robust geochronological studies have been conducted, the gold deposits and proposed fertile granitic intrusions are not the same age (e.g., Goldfarb et al., 2005; Goldfarb and Groves, 2015), and references therein; Vielreicher et al., 2015). Furthermore, the proposed parent granitic rocks have no consistent composition or oxidation state within or between terranes. In some cases, lamprophyres and other more mafic intrusions are close in age to the gold deposits (e.g., Vielreicher et al., 2010), but are volumetrically minor and could not have provided the large volumes of fluids required to form the gold deposits. Although stable isotope data are broadly permissive of a magmatic-hydrothermal fluid, they, combined with the occurrence of some deposits that formed at over 15 km depth and conflicting radiogenic isotope ratios, are indicative of long fluid pathways (e.g., Kontak and Kerrich, 1995; Ridley and Diamond, 2000) that effectively exclude exsolution of ore fluids from granitic intrusions at any reasonable crustal depth. Redox changes, commonly invoked in fluid mixing models (e.g., Walshe et al., 2003; Neumayr et al., 2007) can occur via rock reaction (e.g., Evans et al., 2006) or even during episodic fault rupturing along fluid channelways (e.g., Yamaguchi et al., 2011). It can be concluded that magmatic-hydrothermal processes cannot explain the genesis of individual deposits let alone provide a universal model for orogenic gold formation. Hybrid magmatism with a mixed metasomatized sub-continental lithospheric mantle (SCLM) and crustal source is interpreted to provide the source of fluid and metals for other gold and gold-copper deposit types (e.g., Groves et al., 2010; Mair et al., 2011; Hronsky et al., 2012; Griffin et al., 2013), but cannot have been responsible for formation of economic orogenic gold deposits on the basis of age considerations, lack of volumetrically significant intrusions from this source, and lack of underlying SCLM in some cases (e.g., Goldfarb and Groves, 2015; Groves and Santosh, 2015). Similarly, models involving devolatilization related to emplacement of mantle plumes into the lower crust (e.g., Bierlein and Pisarevsky, 2008; de Boorder, 2012; Webber et al., 2013) lack credible supporting evidence.

This effectively leaves metamorphic models as the only viable possibilities if a universal or near-universal model is sought for the

genesis of orogenic gold deposits. Lateral secretion models (e.g., Boyle, 1979; Saager et al., 1982) are almost certainly invalid due to the limited volume of metamorphic fluid and metals available to produce high-tonnage or high-grade gold deposits (e.g., Glasson and Keays, 1978) even if lateral flow was dominant, which appears unlikely (e.g., Ord and Oliver, 1997). Similarly, a model in which mantle CO₂ was advected through the lower crust to produce granulites and a CO₂-rich pseudo-metamorphic fluid (e.g., Fyon et al., 1984; Cameron, 1988; Santosh and Omori, 1988; Touret and Huizenga, 2012; Fu and Touret, 2014) are not supported by associations between granulites and gold nor by CO₂ contents of fluid inclusions, available carbon isotope data, or theoretical considerations, as originally discussed by Kerrich (1989).

Such considerations have led to general acceptance of a metamorphic model for orogenic gold that promotes metamorphic devolatilization of supracrustal/intrabasinal rocks under greenschist- to amphibolite-facies conditions, with upwards advection of resultant metamorphic fluid and released metals to the site of formation of orogenic gold mineralization at higher crustal levels (e.g., Kerrich and Fyfe, 1981; Phillips and Groves, 1983; Colvine et al., 1984; Goldfarb et al., 1986, 1988; Groves et al., 1987; Cox et al., 1991; Powell et al., 1991; Bierlein and Crowe, 2000; Goldfarb et al., 2001, 2005; Phillips and Powell, 2010; Tomkins, 2010; among many others). This model and its potential limitations are outlined below, as synthesised from the references above and the exhaustive review of Goldfarb and Groves (2015).

3. Supracrustal metamorphic model: strengths and weaknesses

The supracrustal or intrabasinal metamorphic model for orogenic gold deposits is the only one of those discussed above that has the potential to provide a universal model to explain the extraordinary longevity of the deposit type throughout Earth history (Goldfarb et al., 2001). Its strengths are that it requires no specific associations with host rock types, as most lithologies are mineralized in gold provinces globally, nor any association with any specific type of intrusion. It is also consistent with the broadly late-metamorphic and late-deformational timing of gold deposition, and the stable and radiogenic isotope data that are internally ambiguous but collectively suggest long and complex fluid pathways that intersected a variety of rock types (Ridley and Diamond, 2000). The typical low-salinity H₂O-CO₂ (\pm CH₄, N₂) fluid is also that expected from metamorphic devolatilization of supracrustal rocks.

Proponents of the model suggest that auriferous aqueous-carbonic fluids (Phillips and Powell, 2010) are released during greenschist- to amphibolite-facies metamorphism of supracrustal rocks in the mid crust (e.g., Powell et al., 1991). Calculations indicate that up to 5 vol.% of both pelitic and mafic volcanic rocks can be converted to such fluid (e.g., Fyfe et al., 1978; Elmer et al., 2006), providing an adequate fluid flux for even giant deposits (e.g., Phillips and Powell, 2010). Such fluids are envisaged to have migrated to regional fault systems, with which all major deposits are spatially associated, and moved upwards at supralithostatic pressures to deposit gold plus related elements and silica during pressure fluctuations related to seismic events along the fault networks (e.g., Cox et al., 1991). Although such deposition appears to have preferentially occurred in rheologically favourable rocks close to the ductile-brittle transition, deposits formed over a total crustal range from 3 to 15 km and possibly deeper (e.g., Groves, 1993). Both sulfidation reactions and phase separation were involved in gold deposition from a normally reduced, near-neutral fluid carrying gold as a thiosulfide complex. From a tectonic viewpoint, orogenic gold deposits may occur in terranes where

metamorphism was caused by a variety of crustal- to mantle-related processes (Goldfarb et al., 1998). However, gold deposition was broadly coincident with a change in far field stress with resultant transition in deformation from compression to transpression, more rarely transtension, during accretion (e.g., Goldfarb et al., 1988) with concomitant uplift and lowering of lithostatic pressure (e.g., Groves et al., 1987; White et al., 2015).

In a supracrustal metamorphic model, the source rocks for fluid and metals must have changed with time, as discussed in detail by Goldfarb and Groves (2015). In the Phanerozoic, host basins are dominated by sedimentary rocks, commonly turbidite sequences, which host the gold deposits. Mafic volcanic rocks are present in some gold provinces but are not ubiquitous. Hence, metamorphism of a sedimentary source rock is most likely to have produced the ore fluid in the metamorphic model. In contrast, most Precambrian terranes that host orogenic gold deposits, particularly those of Archean age, are dominated by ultramafic to felsic volcanic rock sequences below the crustal level of gold deposition, with ubiquitous mafic rocks the most likely source of auriferous fluid in this model. Specific gold-enriched source rocks have been suggested for both Phanerozoic and Precambrian gold provinces (e.g., Glasson and Keays, 1978; Tomkins, 2010; Large et al., 2011; Steadman et al., 2013), but these are not common to all orogenic gold provinces, are commonly volumetrically insignificant, and cannot be an important factor in any holistic metamorphic model for orogenic gold.

Despite the obvious strengths of the supracrustal metamorphic model, there are some weaknesses, particularly for the Precambrian deposits. These are discussed below, first for Phanerozoic deposits and then for Precambrian examples.

The Phanerozoic deposits fit the model well in that all significant deposits are in greenschist-facies domains (Goldfarb et al., 2005), and research by Pitcairn et al. (2006) on the Otago Schists of New Zealand shows that it is feasible to release significant Au, As, Bi, Sb, Te and W during greenschist- to amphibolite-facies metamorphism of thick turbidite sequences. However, it is still not clear how such fluids can migrate laterally on a kilometre scale into the regional-scale faults that control gold mineralization at the first-order structural scale. Authors such as Ferry (1994) showed that such lateral flow is possible, particularly if induced by external tectonic perturbations such as drainage caused by dilatant fault zones (e.g., Sibson, 1992), but most authors such as Ord and Oliver (1997) suggested that vertical advection would be dominant. Hence there may be a mass balance problem in terms of the amount of gold transported via lateral flow. Furthermore, Ridley (1993) suggested that the high fluid pressure in the regional-scale faults promotes outward flow down pressure gradients into subsidiary faults that host the gold ores. Hence, the mechanism that could promote fluid flow from the metamorphic belt into the regional-scale faults at depth and then back into the rock sequences at higher crustal levels is unclear. The fact that gold mineralization normally postdates regional metamorphism in host sequences, that it is retrogressive (e.g., Wilson et al., 2013), in some instances by several million years (e.g., Perring et al., 1987; Nesbitt, 1991), is another potential problem for both Phanerozoic and Precambrian deposits. This problem is generally overcome by models that suggest that peak metamorphism is attained earlier at deeper crustal levels than at the crustal level of gold deposition (deeper-later model of Stuwe, 1998). However, this does not completely explain why early greenschist-facies metamorphism involves grain-boundary fluid migration with pervasive metamorphic fabrics and assemblages and ubiquitous, totally barren, quartz veins in contrast to the focussed fluid flux that can produce high-grade gold shoots later in the metamorphic history. A possible solution is provided by Goldfarb et al. (1988) who suggested that a change in

far-field stress from compression to transpression might have unlocked metamorphic fluid reservoirs stored in the mid to upper crust as the overlying greenschist-facies domains with their deformational fabrics cooled. This does not completely explain why the fluid was suddenly capable of depositing large quantities of gold when there is no recorded evidence that the earlier fluid was gold-bearing from studies of ubiquitous metamorphic rocks.

As noted above, the metamorphic model for Precambrian deposits has similar problems to its Phanerozoic equivalents. In addition, it is not so clear that ore components can be derived from greenschist- to amphibolite-facies metamorphism of mafic volcanic rocks as it is from turbidites. For example, [Pitcairn et al. \(2015\)](#) demonstrated that, although gold can be liberated in similar concentrations from basalt, in the examples studied from New Zealand, arsenic, the most common associated element in Precambrian orogenic gold deposits, is not liberated during amphibolite-facies metamorphism. A potentially insurmountable problem for the supracrustal model is the occurrence of a significant number of deposits, including the giant Huttu and Kolar deposits in India (e.g., [Sarma et al., 2011](#)), in mid- to upper-amphibolite facies domains that have alteration assemblages that formed under similar P-T conditions to the metamorphosed host rocks ([Colvine et al., 1988](#); [Groves, 1993](#); [Knight et al., 1993](#); [McCuaig et al., 1993](#); [Neumayr et al., 1993](#); [Bloem et al., 1994](#); [Miller and Adams, 2013](#)). Several proponents of the metamorphic model have argued that these deposits formed under lower P-T conditions and were subsequently metamorphosed under amphibolite-facies conditions (e.g., [Tomkins and Mavrogenes, 2002](#); [Tomkins et al., 2004](#); [Tomkins and Grundy, 2009](#); [Phillips and Powell, 2009](#)). However, recent research by [Kolb et al. \(2015\)](#) has demonstrated that a number of these deposits clearly formed under broadly amphibolite-facies conditions, except where they were overprinted during a later, unrelated orogenic event. Hence, the fluid source must have been deeper than the 15 km (possibly up to 20 km) depth of deepest deposit formation. Additional evidence for a deep source, below the supracrustal sequences hosting the gold deposits, is provided by lead isotope evidence from the Neoproterozoic gold province of Western Australia. Importantly, Neoproterozoic greenstone belts can give more-meaningful source area data than for Phanerozoic terranes because much of the lead in the ores may be dominated by the relatively low concentration of lead that is being carried in the ore fluids (e.g., [Goldfarb et al., 2005](#)). [Browning et al. \(1987\)](#) and [McNaughton et al. \(1993\)](#), among others, showed that the lead isotope ratios from the giant Eastern Goldfields Province of the Yilgarn Craton reflect the age and composition of the basement rocks to the supracrustal greenstone belts, implicating a deeper source for the auriferous ore fluids. In confirmation that this is not an isolated instance, orogenic gold deposits throughout the Paleozoic of Ireland ([Standish et al., 2014](#)) are characterized by highly variable lead signatures, reflecting many different lithologies, including the basement.

In summary, although the supracrustal metamorphic model satisfies the majority of constraints from geological, geochronological, geochemical, isotopic and fluid inclusion data, there are a number of weaknesses in terms of a unified model. The Precambrian examples provide the greatest problems, with evidence that ore fluids were derived from below 15 km depth and carry components that must have been derived from the basement to the gold-hosting greenstone belts. Furthermore, there is doubt that all metal components can be derived from greenschist- to amphibolite-facies metamorphism of a basaltic rock, the only volumetrically viable source in the greenstone belts, in contrast to evidence suggesting that such components can be derived from sedimentary sources. There is also the problem of definition of the precise mechanism by which ore fluid migrated laterally on a kilometre scale into the regional-scale faults that clearly control the

location of gold districts and provinces. Finally, the model does not adequately explain the conjunction of apparent late- to post-metamorphic timing in host sequences precisely at the time that a change in far-field stresses promoted a change from compression, represented by the metamorphic fabrics in the host rocks, to transpression or transtension, demonstrated by the geometry of the orogenic gold ore bodies.

The ore fluid is clearly a metamorphic fluid but the evidence above suggests a source below the supracrustal sequences that host the gold deposits. As argued above, this cannot be the lower crust, nor the metasomatized lithosphere, nor a mantle plume and associated granulitization. Similarly, the ore fluid is unlikely to be exactly the same fluid that caused regional metamorphism of supracrustal rocks and related quartz veins with no evidence of any gold enrichment. This specific metamorphic ore fluid appears to have been liberated at a unique time in the orogenic cycle during a change in far-field stress. As orogenic gold deposits are inevitably formed in accretionary or, less commonly, collisional tectonic environments related to subduction ([Goldfarb et al., 2001, 2005](#)), and not in other types of metamorphic belts, this suggests a fundamental relationship to a change in plate motion.

In seeking an explanation and a unified model, it is important to view those deposits that cannot have formed from a metamorphic fluid derived from within the host supracrustal sequences because they had been metamorphosed to at least the amphibolite-facies hundreds to thousands of million years previously. Such Tertiary deposits occur along the Megashear Zone in the Proterozoic–Phanerozoic terranes of northern Mexico ([Iriando, 2001](#); [Goldfarb et al., 2007](#)) and Cretaceous deposits occur in the Archean–Proterozoic terranes of the North China block. The giant Jiaodong gold province of the latter (e.g., [Wang et al., 1998](#); [Goldfarb et al., 2007](#); [Goldfarb and Santosh, 2014](#); [Yang et al., 2014](#)) is discussed below in an attempt to solve the problems discussed above and seek to develop a unified model that can explain the origin of all orogenic gold deposits of all ages.

4. The giant Jiaodong gold province: the exception or the key to a unified model

The giant Jiaodong gold province in the eastern half of the North China block ([Li et al., 2015a](#); [Song et al., 2015](#); [Yang and Santosh, 2015](#); [Yang et al., 2016a](#)) represents a region of major lithospheric erosion of originally thick buoyant Archean sub-continental lithospheric mantle or SCLM ([Griffin et al., 1998](#); [Santosh, 2010](#)), caused by anomalously complex Mesozoic slab subduction from the north, south, and east. This led to slab devolatilization, subsequent melting, and voluminous granitic magmatism ([Windley et al., 2010](#)). The associated Yanshanian orogeny, that occurred within the decratonized North China block, was typified by basement uplift, regional extension, ca. 165–90 Ma granite intrusion, and ca. 130–120 Ma gold formation within the eastern margins of this highly modified cratonic basement ([Goldfarb and Santosh, 2014](#); [Yang and Santosh, 2015](#)). The structural control and more protracted period of gold mineralization argue against a magmatic-hydrothermal fluid model, and the Precambrian high-grade metamorphism of the basement rocks clearly invalidates a supracrustal metamorphic-devolatilization fluid model for the gold event. Despite this, the widespread gold episode correlates with changing far-field stresses and plate reorganizations as interpreted for other orogenic gold provinces.

For these reasons, the Jiaodong deposits have generally been classified as orogenic gold deposits ([Wang et al., 1998](#); [Goldfarb et al., 2001, 2005](#); [Yang et al., 2016a](#)). Although they are commonly hosted by older granites, they show no close spatial relationship to granitic intrusions of the same age, nor evidence of metal zonation related to

thermal gradients around hot intrusions. However, they do show a clear structural control along regional faults, and ore and wallrock-alteration mineralogy, fluid inclusion compositions and stable isotope chemistry are similar to more typical orogenic gold deposits, particularly of epizonal type (Yang et al., 2016b).

As noted above, however, in contrast to the normal situation where orogenic gold deposits formed within 50–200 my of the deposition of their host supracrustal sequence (Goldfarb et al., 2001), the Jiaodong deposits formed at ca. 126–120 Ma, some 2 Ga after the oldest host rocks in the North China block were deposited and experienced high P-T metamorphism (Yang et al., 2014; see review in Goldfarb and Santosh (2014)). Importantly, prograde metamorphism of supracrustal host rocks cannot have provided auriferous fluids and a sub-crustal source must have provided the fluid and metals that are interpreted to have advected up the Tan Lu and other crustal-scale fault systems (Fig. 2) in the province (Goldfarb et al., 2007). Deep crustal-basement sources are highly unlikely, as discussed more generally above, implicating a sub-crustal source. Although the metasomatized SCLM may be gold-enriched (e.g., Griffin et al., 2013) and may be the source of magmas and fluids for some gold and gold-copper deposits (e.g., Hronsky et al., 2012), this magmatic-hydrothermal model can effectively be ruled out for orogenic gold formation generally (Groves and Santosh, 2015), as discussed above. However, direct devolatilization of an overlying frozen mantle wedge cannot be completely ruled out as a fluid source under specific conditions (e.g., Wyman et al., 2008).

This leaves the subducted oceanic crust and overlying sediment wedge as the most viable fluid and metal source. Goldfarb and Santosh (2014) and Goldfarb and Groves (2015) evaluated how these could have provided the source of the Jiaodong ore fluids, with Fig. 3 adapted from their model. Basically, devolatilization of a subducted slab can result in extensive upward fluid-flux along slab-mantle boundaries (e.g., Sibson, 2004; Peacock et al., 2011) into fore-arc or accreting terrane margins. Such metamorphic fluid release occurs when the base of the fore-arc mantle wedge becomes fully hydrated (Katayama et al., 2012). At this stage, the oceanic slab will devolatilize, together with its overlying pyrite-bearing oceanic sediment wedge. The latter is important as most proponents of specific source rocks for formation of orogenic gold




Figure 3. Schematic illustration of the slab devolatilization model for the formation of orogenic gold deposits of the giant Jiaodong gold province as described in the text. Adapted from Goldfarb and Santosh (2014) and Goldfarb and Groves (2015).

deposits stress the importance of gold-enriched pyrite in sediments or sedimentary rocks, as gold and related elements can be released to the fluid via breakdown of pyrite to pyrrhotite (e.g., Large et al., 2009, 2011; Steadman et al., 2013). The highly sheared serpentized layer at the bottom of the corner of the mantle wedge may provide a particularly permeable zone for slab dewatering at slab depths of less than 100 km and temperatures of 650 °C (Kawano et al., 2011). Over-pressured fluids (e.g., Sibson, 2013) could then migrate up-dip, channelling into crustal-scale fault zones at higher crustal levels to eventually deposit orogenic gold deposits at even shallower levels in lower-order structures (e.g., Breeding and Ague, 2002; Hyndaman et al., 2015).

Goldfarb and Santosh (2014) applied this model to the giant Jiaodong orogenic gold province, suggesting that the Tan Lu fault system carried the auriferous fluid derived from slab devolatilization up to higher crustal levels to deposit gold in the numerous deposits in the province. An important factor in such a model is the trigger to cause fluid release in the slab and sediment wedge. As discussed by Seno and Kirby (2014), such a trigger might be the end of subduction or stalling of the slab during subduction, which could




Figure 2. Tectonic framework of the North China Craton showing the distribution of gold deposits. The three major Paleoproterozoic sutures (Inner Mongolia Suture Zone, Trans-North China Orogen and Jiao-Liao-Ji Belt) along which the crustal blocks amalgamated at the final stage of cratonization during late Paleoproterozoic are also shown. Adapted from Groves and Santosh (2014).

result in a change of stress regime as plates were reorganised with subsequent switchover from compression to transpression. Wyman et al. (2008) and Wyman and Kerrich (2010) further suggested that such a devolatilization process would be favoured by flat subduction and could have extended back to the Archean.

That such fluids can be transported from the mantle to crustal levels in crustal-scale fault zones is shown by radiogenic isotope, halogen and noble gas data of the San Andreas fault system (Kennedy and van Soest, 2007; Pili et al., 2011) and the Karakorum fault zone (Klemperer et al., 2013).

Although it must be stressed that the slab devolatilization model is only a hypothetical one, because all radiogenic and stable isotope ratios and other geochemical data are equivocal for any model for the genesis of orogenic gold deposits, it is the only pragmatic model that can satisfy the available geological and geochronological constraints for Jiaodong.

Goldfarb and Groves (2015) went further in suggesting that the slab-source model is the only reasonable genetic model for the gold deposits in the potentially giant Qinling gold province on the southern margin of the North China block (e.g., Chen et al., 2008), the region marking the closure of the northernmost paleo-Tethys sea and the tectonic suturing of the Yangtze and North China Cratons (Li et al., 2015b). Goldfarb et al. (2007) also noted that the Tertiary gold deposits in northwestern Mexico and southernmost Arizona are hosted in reactivated high-grade Proterozoic basement in extensional structures within metamorphic core complexes in a similar scenario to the deposits in the North China block.

The question then remains whether the giant Jiaodong gold province, and the others with similar features mentioned above, are highly anomalous members of the orogenic gold clan or are the key to understanding all orogenic gold deposits in terranes of all ages. Certainly, the Goldfarb and Santosh (2014) Jiaodong model can explain the ubiquitous worldwide relationship to subduction and generation of new crust, the late-metamorphic timing of orogenic gold mineralization, and the temporal coincidence of auriferous fluid release and transition from compression to transpression if a stalled slab resulted in a change in far-field stress. It can also explain the occurrence of Precambrian deposits at crustal depths below those inferred for the supracrustal metamorphic model and the lead isotope evidence that fluids interacted with basement. It also obviates the somewhat problematic issue of extensive lateral fluid flow, as the crustal-scale faults extend to the Moho and slab-related metamorphic fluids could be channelled directly into them with subsequent upward flow. It also elegantly explains the conflicting isotopic and halogen and rare gas evidence for upper to lower crustal and mantle components. Finally, it overcomes the problem of implication of different source rocks with time as the most favoured source rocks, gold-enriched pyrite-rich marine sediments, would have been available in sediment wedges above down-going slabs throughout Earth history. Metamorphic heating would have consistently transformed pyrite to pyrrhotite, releasing gold and related elements such as Ag, As, Bi, Sb, Te and W into a sulfur-bearing ore fluid. The higher CO₂ content of the Archean ore fluids could also be explained by a greater degree of carbonation in Archean oceanic rocks due to a combination of more-susceptible high-MgO basalts and lack of CO₂ sinks. Santosh and Omori (2008) and Santosh et al. (2009) evaluated the history of CO₂ circulation through time and proposed schematic models for the Archean and post-Archean scenarios which can further enhance a model for greater degrees of carbonation of gold and fluid source rocks in the Archean (Fig. 4). The Mid Oceanic Ridge basalt, carbonated at spreading axes and during lateral transport associated with seafloor spreading, is eventually decarbonated during subduction at convergent plate boundaries. They speculated that the incorporation of carbonate into the




Figure 4. Schematic illustrations showing CO₂ circulation through time for Archean and post-Archean scenarios (after Santosh and Omori, 2008; Santosh et al., 2009).

subcontinental lithosphere by subduction was probably initiated by at least 4 Ga. Oceanic carbonates infiltrate into the deep mantle domains during subduction, with carbonated mantle underlying ancient (>2.0 Ga old) continents. The fluids released from these domains would have advected up the crustal-scale faults under steady-state creep and then migrated episodically along pressure and permeability gradients during earthquake activity into lower-order structures and/or hydraulically-fractured rock bodies above the ductile-brittle transition (e.g., Cox et al., 2001).




Figure 5. Schematic representation of a permissive scenario for all orogenic gold deposits, including those in high-grade metamorphic rocks, involving a subcrustal fluid and metal source from slab devolatilization. (a) Where these overpressured slab-derived fluids intersect deep-crustal faults, they advect upwards to form orogenic gold deposits in second-order structures or hydraulically-fractured rock bodies: based on Archean lithostratigraphic controls on ore bodies. (b) Fluids released during devolatilization of the subducting slab and associated sulfidic sediments at temperatures below 650 °C and depths of 100 km may either fertilize the overlying mantle wedge or, particularly once the wedge is fully hydrated, travel up-dip along the interface between the slab and the overlying serpentinized wedge or base of the lithosphere. Figure adapted from Goldfarb and Groves (2015). Detailed models are presented in Wyman et al. (2008).

Thus, the slab devolatilization model, shown schematically in Fig. 5, can potentially explain the conjunction of parameters that consistently characterize the orogenic gold deposit clan. Like all models, they suffer for a lack of complete understanding of the precise processes operating at depth in the system, and in this specific case, the lack of knowledge of subduction zone geometry at the time of gold mineralization and whether analogies to fluid migration into modern fore-arcs can be made to accreted terranes containing back-arc components at the time of slab devolatilization. However, the model is considered here to be the most consistent if the fundamental concept of a unified model for orogenic gold systems, such as those for other coherent mineral deposit groups, is valid.

5. Concluding statement

There are no modern examples of orogenic gold deposits, with the youngest well-documented, but uneconomic, orogenic gold mineralization forming about 12–15 Ma ago in the European Alps, and the most recent significant orogenic gold deposit forming about 50 Ma ago (Goldfarb et al., 2001). They also form over a crustal depth range unparalleled by any other gold deposit group with the possible exception of iron-oxide copper-gold deposits (e.g., Groves et al., 2010), in very complex tectonic environments where various mantle and crustal processes could be involved in crustal heating and fluid release. Therefore, it is to be expected that their origin is controversial and that research tools such as fluid inclusions, stable and radiometric isotope ratios and other geochemical methodologies should prove equivocal. Such genetic controversy has existed for a century or so for gold-only lode-gold deposits, and continues today despite their recognition as a coherent orogenic gold deposit group by Groves et al. (1998).

If the deposit group has a coherent set of critical features, like all other deposit groups, it should share a specific origin and relate to a unified minerals-system model. Based on consistent relationships and geochronological constraints, two variants of a metamorphic model are the only possibilities of providing such a unified model: (1) a model involving devolatilization of deeper supracrustal sequences underlying the host rocks to the deposits, and (2) a model involving devolatilization of a subducted slab and overlying sediment wedge.

The supracrustal metamorphic model requires that source rocks varied from mafic volcanic rocks to sedimentary rocks from the Precambrian to Phanerozoic, that auriferous metamorphic fluid derived at greenschist- to amphibolite-facies P-T conditions was expelled to higher crustal levels and resulted in mineralization in rocks that were already metamorphosed in the same event, and that significant lateral flow of such fluid towards crustal-scale faults was possible. In such a model, the Precambrian gold deposits deposited under amphibolite-facies conditions and the young deposits formed in much older, previously metamorphosed host rocks, such as in the giant Jiaodong gold province, have to be considered anomalous exceptions to the model. It is thus not a unified model for all orogenic gold deposits of all ages.

In contrast, the slab devolatilization hypothesis, developed for the giant Jiaodong gold province, has the potential to be a unified model (Fig. 5) that can incorporate all orogenic gold deposits including the high-PT Precambrian deposits. It can explain why there is a consistent connection between post-peak metamorphic auriferous-fluid advection and change in tectonic regime due to stalling of a subducted slab, how over-pressured fluid can migrate directly upwards into crustal-scale fault zones and then down hydraulic gradients into gold depositional sites at the ductile-brittle transition, and involves a common fluid source, the slab and overlying pyritic sediment wedge, throughout geological history. It

is truly a holistic model, whose only uncertainty, shared with all models, is the precise processes operating at depth: in this case exactly how fluid is channelled along the slab-mantle boundary and how this fluid can migrate ocean-wards to the accreted terranes that typically host these orogenic gold deposits.

In conclusion, it is believed that the slab devolatilization model is the model that explains most of the tectonic- to deposit-scale features of orogenic gold deposits throughout Earth history and requires less special pleading of specific explanations for individual anomalous and controversial examples. In this view, the giant Jiaodong gold province is the key to unlocking a unified minerals-system model, not the exception to a supracrustal metamorphic model.

Acknowledgements

We are indebted to many colleagues over many years who have contributed to our knowledge of gold mineralization and associated tectonic environments and processes. We are particularly indebted to Rich Goldfarb of the USGS for his invaluable contributions including the enlightenment he has provided on the Jiaodong gold province. He also made available a paper in *Lithos* by Goldfarb and Groves (2015) which we have unashamedly used as a source of information and references on orogenic gold. He is only not a co-author of the paper because we differ in our view of the significance of Jiaodong to holistic models for orogenic gold deposits.

References

- Baker, T., 2002. Emplacement depth and CO₂-rich fluid inclusions in intrusion-related gold deposits. *Economic Geology* 97, 1109–1115.
- Bath, A.B., Walshe, J.L., Cloutier, J., Verrall, M., Cleverley, J.S., Pownceby, M.I., Macrae, C.M., Wilson, N.C., Tunjic, J., Nortje, G.S., Robinson, P., 2013. Biotite and apatite as tools for tracking pathways of oxidized fluids in the Archean east repulse gold deposit, Australia. *Economic Geology* 108, 667–690.
- Bierlein, F.P., Crowe, D.E., 2000. Phanerozoic orogenic lode gold deposits. *Reviews in Economic Geology* 13, 103–139.
- Bierlein, F.P., Groves, D.I., Goldfarb, R.J., Dube, B., 2006. Lithospheric controls on the formation of giant orogenic gold deposits. *Mineralium Deposita* 40, 874–886.
- Bierlein, F.P., Pisarevsky, S., 2008. Plume-related oceanic plateaus as a potential source of gold mineralization. *Economic Geology* 103, 425–430.
- Bloem, E.J.M., Dalstra, H.J., Groves, D.I., Ridley, J.R., 1994. Metamorphic and structural setting of Archean amphibolite-hosted gold deposits near Southern Cross, Southern Cross Province, Yilgarn Block, Western Australia. *Ore Geology Reviews* 9, 183–208.
- Boyle, R.W., 1979. The geochemistry of gold and its deposits. *Geological Survey of Canada Bulletin* 280, 584.
- Breeding, C.M., Ague, J.J., 2002. Slab-derived fluids and quartz-vein formation in an accretionary prism, Otago Schist, New Zealand. *Geology* 30, 499–502.
- Browning, P., Groves, D.I., Blockley, J.G., Rosman, K.J.R., 1987. Lead isotope constraints on the age and source of gold mineralization in the Archean Yilgarn Block, Western Australia. *Economic Geology* 82, 971–986.
- Cameron, E.M., 1988. Archean gold—relation to granulite formation and redox zoning in the crust. *Geology* 16, 109–112.
- Chen, Y.-J., Pirajno, F., Qi, J.-P., 2008. The Shanggong gold deposit, Eastern Qinling Orogen, China— isotope geochemistry and implication for ore genesis. *Journal of Asian Earth Sciences* 33, 252–266.
- Cline, J.S., Hofstra, A.H., Muntean, J.L., Tosdal, R.M., Hickey, K.A., 2005. Carlin-type gold deposits in Nevada: critical geologic characteristics and viable models. *Economic Geology 100th Anniversary Volume*, 451–484.
- Colvine, A.C., Andrews, A.J., Cherry, M.E., Durocher, M.E., Fyon, J.A., Lavigne, M.J., Macdonald, A.J., Marmont, S., Poulsen, K.H., Springer, J.S., Troop, D.G., 1984. An Integrated Model for the Origin of Archean Lode-gold Deposits. Ontario Geological Survey Open-File Report 5524, p. 98.
- Colvine, A.C., Fyon, J.A., Heather, K.B., Marmont, S., Smith, P.M., Troop, D.G., 1988. Archean lode gold deposits in Ontario. Ontario Geological Survey Miscellaneous Paper 139, 136.
- Cox, S.F., Etheridge, M.A., Cas, R.A.F., Clifford, B.A., 1991. Deformational style of the Castlemaine area, Bendigo-Ballarat Zone: implications for evolution of the crustal structure across southeast Australia. *Australian Journal of Earth Sciences* 38, 151–170.
- Cox, S.F., Knackstedt, M.A., Braun, J., 2001. Principles of structural control on permeability and fluid flow in hydrothermal systems. *Society of Economic Geology Reviews* 14, 1–24.

- de Boorder, H., 2012. Spatial and temporal distribution of the orogenic gold deposits in the Late Palaeozoic Variscides and Southern Tianshan—How orogenic are they? *Ore Geology Reviews* 46, 1–31.
- Elmer, F.L., White, R.W., Powell, R., 2006. Devolatilisation of metabasic rocks during greenschist-amphibolite facies metamorphism. *Journal of Metamorphic Geology* 24, 497–513.
- Evans, K.A., Phillips, G.N., Powell, R., 2006. Rock-buffering of auriferous fluids in altered rocks associated with the Golden Mile-style mineralization, Kalgoorlie gold field, Western Australia. *Economic Geology* 805–817.
- Ferry, J.M., 1994. Overview of the petrological record of fluid flow during regional metamorphism in northern New England. *American Journal of Science* 294, 905–988.
- Fu, B., Touret, J.L.R., 2014. From granulite fluids to quartz-carbonate megashear zones: the gold rush. *Geoscience Frontiers* 5, 747–758.
- Fyfe, W.S., Price, N.J., Thompson, A.B., 1978. *Fluids in the Earth's Crust*. Elsevier, Amsterdam, p. 383.
- Fyon, J.A., Schwarcz, H.P., Crocker, J.H., 1984. Carbonatization and gold mineralization in the Timmins area, Abitibi greenstone belt: genetic links with Archean mantle CO₂-degassing and lower crust granulitization, vol. 9. Geological Association of Canada Program with Abstracts, p. 65.
- Gebre-Mariam, M., Hagemann, S.G., Groves, D.I., 1995. A classification scheme for epigenetic Archean lode-gold deposits. *Mineralium Deposita* 30, 408–410.
- Glasson, M.J., Keays, R.R., 1978. Gold mobilization during cleavage development in sedimentary rocks from the auriferous slate belt of central Victoria, Australia; some important boundary constraints. *Economic Geology* 73, 496–511.
- Goldfarb, R.J., Baker, T., Dubé, B., Groves, D.I., Hart, C.J.R., Gosselin, P., 2005. Distribution, character, and genesis of gold deposits in metamorphic terranes. *Economic Geology* 100th Anniversary Volume, 407–450.
- Goldfarb, R.J., Groves, D.I., 2015. Orogenic gold: common or evolving fluid and metal sources through time. *Lithos* 233, 2–26.
- Goldfarb, R.J., Groves, D.I., Gardoll, S., 2001. Orogenic gold and geologic time: a global synthesis. *Ore Geology Reviews* 18, 1–75.
- Goldfarb, R.J., Hart, C.J.R., Davis, G., Groves, D.I., 2007. East Asian gold: deciphering the anomaly of Phanerozoic gold in Precambrian cratons. *Economic Geology* 102, 341–346.
- Goldfarb, R.J., Hart, C.J.R., Marsh, E.E., 2008. Orogenic gold and evolution of the Cordilleran orogen. In: Spencer, J.E., Tittley, S.R. (Eds.), *Circum-Pacific Tectonics, Geologic Evolution, and Ore Deposits*, vol. 22. Tucson Arizona Geological Society Digest, pp. 311–323.
- Goldfarb, R.J., Leach, D.L., Miller, M.L., Pickthorn, W.J., 1986. Geology, Metamorphic Setting, and Genetic Constraints of Epigenetic Lode-gold Mineralization within the Cretaceous Valdez Group, South-central Alaska. *Geological Association of Canada Special Paper* 32, pp. 87–105.
- Goldfarb, R.J., Leach, D.L., Pickthorn, W.J., Paterson, C.J., 1988. Origin of lode-gold deposits of the Juneau gold deposit, southeast Alaska. *Geology* 16, 440–443.
- Goldfarb, R.J., Phillips, G.N., Nokleberg, W.J., 1998. Tectonic setting of synorogenic gold deposits of the Pacific Rim. *Ore Geology Reviews* 13, 185–218.
- Goldfarb, R.J., Santosh, M., 2014. The dilemma of the Jiaodong gold deposits: are they unique? *Geoscience Frontiers* 5, 139–153.
- Goldfarb, R.J., Taylor, R.D., Collins, G.S., Goryachev, N.A., Orlandini, O.F., 2014. Phanerozoic continental growth and gold metallogeny of Asia. *Gondwana Research* 25, 48–102.
- Griffin, W.L., Begg, G.C., O'Reilly, S.Y., 2013. Continental root control on the genesis of magmatic ore deposits. *Nature Geoscience* 6, 905–910.
- Griffin, W.L., Zhang, A., O'Reilly, S.Y., 1998. Phanerozoic evolution of the lithosphere beneath the Sino-Korean craton. In: Folwer, M.F.J. (Ed.), *Mantle Dynamics and Plate Interactions in East Asia*. American Geophysical Union Geodynamics Series 27, pp. 107–126.
- Groves, D.I., 1993. The crustal continuum model for late-Archaean lode gold deposits of the Yilgarn block, Western Australia. *Mineralium Deposita* 28, 366–374.
- Groves, D.I., Bierlein, F.P., Meinert, L.A., Hitzman, M.W., 2010. Iron oxide copper-gold (IOCG) deposits through Earth history: implications for origin, lithospheric setting and distinction from other epigenetic iron-oxide deposits. *Economic Geology* 105, 641–654.
- Groves, D.I., Goldfarb, R.J., Gebre-Mariam, M., Hagemann, S.G., Robert, F., 1998. Orogenic gold deposits: a proposed classification in the context of their crustal distribution and relationship to other gold deposit types. *Ore Geology Reviews* 13, 7–27.
- Groves, D.I., Goldfarb, R.J., Santosh, M., 2016. The conjunction of factors that lead to the formation of giant gold provinces and deposits in non-arc settings. *Geoscience Frontiers* 7 (3), 303–314.
- Groves, D.I., Santosh, M., 2015. Province-scale commonalities of some world-class gold deposits: implications for mineral exploration. *Geoscience Frontiers* 6, 389–399.
- Groves, D.I., Phillips, G.N., Ho, S.E., Houstoun, S.M., Standing, C.A., 1987. Craton-scale distribution of Archean greenstone gold deposits—predictive capacity of the metamorphic model. *Economic Geology* 82, 2045–2058.
- Hall, G., Wall, V., 2007. Geology works—the use of regional geological maps in exploration. In: Milereit, B. (Ed.), *Proceedings of Exploration 07: Fifth Decennial Conference on Mineral Exploration*, pp. 51–60.
- Hart, C.J.R., McCoy, D., Goldfarb, R.J., Smith, M., Roberts, P., Hulstein, R., Bakke, A.A., Bundtzen, T.K., 2002. *Geology, exploration and discovery in the Tintina gold province, Alaska and Yukon*. Society of Economic Geologists Special Publication 9, 241–274.
- Helt, K.M., Williams-Jones, A.E., Clark, J.R., Wing, B.A., Wares, R.P., 2014. Constraints on the genesis of the Archean oxidized, intrusion-related Canadian Malartic gold deposit, Quebec, Canada. *Economic Geology* 109, 713–735.
- Hronsky, J.M.A., Groves, D.I., Loucks, R.R., Begg, G.C., 2012. A unified model for gold mineralisation in accretionary orogens and implications for regional-scale exploration targeting methods. *Mineralium Deposita* 47, 339–358.
- Hughes, M.J., Phillips, G.N., Gregory, L.M., 1997. *Mineralogical Domains in the Victorian Gold Province, Maldon, and Carlin-style Potential*. Australasian Institute of Mining and Metallurgy, Annual Conference, Melbourne, pp. 215–227.
- Hutchinson, R.W., 1987. Metallogeny of Precambrian gold deposits: space and time relationships. *Economic Geology* 90, 1918–1933.
- Hutchinson, R.W., Burlington, J.L., 1984. Some broad characteristics of greenstone belt gold lodes. In: Foster, R.P. (Ed.), *Gold'82—the Geology, Geochemistry and Genesis of Gold Deposits*. Balkema, Rotterdam, pp. 339–372.
- Hyndman, R.D., McCrory, P.A., Wech, A., Kao, H., Ague, J., 2015. Cascadia subducting plate fluids channeled to forearc mantle corner: ETS and silica deposition. *Journal of Geophysical Research*. <http://dx.doi.org/10.1002/2015JB011920>.
- Iriondo, A., 2001. Proterozoic Basements and Their Laramide Juxtaposition in NW Sonora Mexico—tectonic Constraints on the SW Margin of Laurentia. University of Colorado, Boulder, p. 222 (Unpublished Ph.D. dissertation).
- Katayama, I., Terada, T., Okazaki, K., Tanikawa, W., 2012. Episodic tremor and slow slip potentially linked to permeability contrasts at the Moho. *Nature Geoscience* 5, 731–734.
- Kawano, S., Katayama, I., Okazaki, K., 2011. Permeability anisotropy of serpentinite and fluid pathways in a subduction zone. *Geology* 39, 939–942.
- Kennedy, B.M., van Soest, M.C., 2007. Flow of mantle fluid through the ductile lower crust: helium isotope trends. *Science* 318, 1433–1436.
- Kerrich, R., 1983. Geochemistry of gold deposits in the Abitibi Greenstone Belt, Special Volume 27. Canadian Institute of Mining and Metallurgy, p. 75.
- Kerrich, R., 1989. Archean gold: relation to granulite formation or felsic intrusions? *Geology* 17, 1011–1015.
- Kerrich, R., Fyfe, W.S., 1981. The gold-carbonate association: source of CO₂ and CO₂ fixation reactions in Archean lode deposits. *Chemical Geology* 33, 265–294.
- Klemperer, S.L., Kennedy, B.M., Sastry, S.R., Makovsky, Y., Harinarayana, T., Leech, M., 2013. Mantle fluids in the Karakoram fault: helium isotope evidence. *Earth and Planetary Science Letters* 366, 59–70.
- Knight, J.T., Groves, D.I., Ridley, J.R., 1993. The Coolgardie Goldfield, Western Australia: district-scale controls on an Archean gold camp in an amphibolite facies terrane. *Mineralium Deposita* 28, 436–456.
- Kolb, J., Dziggel, A., Bagas, L., 2015. Hypozonal lode gold deposits: a genetic concept based on a review of the New Consort, Hutti, Hira Buddini, Navachab, Nevoria, and the Granites deposits. *Precambrian Research* 262, 20–44.
- Kontak, D.J., Kerrich, R., 1995. Geological and geochemical studies of a metabasite-hosted lode gold deposit: the Beaver Dam deposit, Nova Scotia: II. Isotopic studies. *Economic Geology* 90, 885–901.
- Lang, J.R., Baker, T., Hart, C.J.R., Mortensen, J.K., 2000. An exploration model for intrusion-related gold systems. *Society of Economic Geology Newsletter* 40 (1), 6–15.
- Large, R.R., Bull, S.W., Maslennikov, V.V., 2011. A carbonaceous sedimentary source-rock model for Carlin-type and orogenic gold deposits. *Economic Geology* 106, 331–358.
- Large, R.R., Danyushevsky, L.V., Hollit, C., Maslennikov, V., Meffre, S., Gilbert, S., Bull, S., Scott, R., Emsbo, P., Thomas, H., Foster, J., 2009. Gold and trace element zonation in pyrite using a laser imaging technique: implications for the timing of gold in orogenic and Carlin-style sediment-hosted deposits. *Economic Geology* 104, 635–668.
- Li, L., Santosh, M., Li, S.-R., 2015a. The 'Jiaodong type' gold deposits: characteristics, origin and prospecting. *Ore Geology Reviews* 65, 589–611.
- Li, N., Chen, Y.J., Santosh, M., Pirajno, F., 2015b. Compositional polarity of triassic granitoids in the Qinling Orogen, China: implication for termination of the northernmost paleo-Tethys. *Gondwana Research* 27, 244–257.
- Mair, J.L., Farmer, G.L., Groves, D.I., Hart, C.J.R., Goldfarb, R.J., 2011. Petrogenesis of postcollisional magmatism at Scheelite Dome, Yukon, Canada: evidence for a lithospheric mantle source for magmas associated with intrusion-related gold systems. *Economic Geology* 106, 451–480.
- McCuag, T.C., Kerrich, R., Groves, D.I., Archer, N., 1993. The nature and dimensions of regional and local gold-related hydrothermal alteration in tholeiitic metabasalts in the Norseman Goldfields: the missing link in a crustal continuum of gold deposits? *Mineralium Deposita* 28, 420–435.
- McNaughton, N.J., Groves, D.I., Witt, W.K., 1993. The source of lead in Archean lode-gold deposits of the Menzies-Kalgoorlie-Kambalda region, Yilgarn Block, Western Australia. *Mineralium Deposita* 28, 495–502.
- Miller, J.M., Adams, G., 2013. The amphibolite-facies Lindsays deposit, Coolgardie Goldfield, Yilgarn Craton, Western Australia—a type example of the crustal continuum model? In: 12th Biennial SGA Meeting, Proceedings, vol. 3. Sweden, Uppsala, pp. 1156–1159.
- Mueller, A.G., 1992. Petrogenesis of amphibole-biotite-calcite-plagioclase alteration and laminated gold-silver quartz veins in four Archean shear zones of the Norseman district, Western Australia. *Canadian Journal of Earth Science* 29, 388–417.
- Nesbitt, B.E., 1991. Phanerozoic gold deposits in tectonically active continental margins. In: Foster, R.P. (Ed.), *Gold Metallogeny and Exploration*. Blackie and Sons Ltd., Glasgow, pp. 104–132.
- Neumayr, P., Cabri, L.J., Groves, D.I., Mikucki, E.J., Jackman, J.A., 1993. The mineralogical distribution of gold and relative timing of gold mineralization in two

- Archean settings of high metamorphic grade in Australia. *Canadian Mineralogist* 31, 711–725.
- Neumayr, P., Walshe, J., Hagemann, S., Petersen, K., Roache, A., Frikken, P., Horn, L., Halley, S., 2007. Oxidized and reduced mineral assemblages in greenstone belt rocks of the St. Ives gold camp, Western Australia: vectors to high-grade ore bodies in Archean gold deposits. *Mineralium Deposita* 43, 363–371.
- Ord, A., Oliver, N.M.S., 1997. Mechanical controls on fluid flow during regional metamorphism: some numerical models. *Journal of Metamorphic Geology* 15, 345–359.
- Peacock, S.M., Christensen, N.I., Bostock, M.G., Audet, P., 2011. High pore pressures and porosity at 35 km depth in the Cascadia subduction zone. *Geology* 39, 471–474.
- Perring, C.S., Groves, D.I., Ho, S.E., 1987. Constraints on the source of auriferous fluids for Archean gold deposits. In: Ho, S.E., Groves, D.I. (Eds.), *Recent Advances in Understanding Precambrian Gold Deposits*, Publication 11, Geology Department and University Extension. University of Western Australia, pp. 287–306.
- Phillips, G.N., Groves, D.I., 1983. The nature of Archean gold-bearing fluids as deduced from gold deposits of Western Australia. *Journal of the Geological Society of Australia* 30, 25–39.
- Phillips, G.N., Groves, D.I., Martyn, J.E., 1984. An epigenetic origin for banded iron-formation-hosted gold deposits. *Economic Geology* 79, 162–171.
- Phillips, G.N., Powell, R., 2009. Formation of gold deposits—review and evaluation of the continuum model. *Earth-Science Reviews* 94, 1–21.
- Phillips, G.N., Powell, R., 2010. Formation of gold deposits—a metamorphic devolatilization model. *Journal of Metamorphic Geology* 28, 689–718.
- Phillips, G.N., Powell, R., 2015. A practical classification of gold deposits, with a theoretical basis. *Ore Geology Reviews* 65, 568–573.
- Pili, E., Kennedy, B.M., Conrad, M.E., Gatiér, J.-P., 2011. Isotopic evidence for the infiltration of mantle and metamorphic CO₂-H₂O fluids from below in faulted rocks from the San Andreas fault system. *Chemical Geology* 281, 242–252.
- Pitcairn, I.K., Craw, D., Teagle, D.A.H., 2015. Metabasalts as sources of metals in orogenic gold deposits. *Mineralium Deposita* 50, 373–390.
- Pitcairn, I.K., Teagle, D.A.H., Craw, D., Olivo, G.R., Kerrich, R., Brewer, T.S., 2006. Sources of metals and fluids in orogenic gold deposits: insights from the Otago and Alpine Schists, New Zealand. *Economic Geology* 101, 1525–1546.
- Powell, R., Will, T., Phillips, G., 1991. Metamorphism in Archean greenstone belts: calculated fluid compositions and implications for gold mineralization. *Journal of Metamorphic Geology* 9, 141–150.
- Ridley, J.R., 1993. The relations between mean rock stress and fluid flow in the crust: with reference to vein- and lode-style gold deposits. *Ore Geology Reviews* 8, 23–37.
- Ridley, J.R., Diamond, L.W., 2000. Fluid chemistry of orogenic lode gold deposits and implications for genetic models. *Reviews in Economic Geology* 13, 141–162.
- Saager, R., Meyer, M., Muff, R., 1982. Gold distribution in supracrustal rocks from Archean greenstone belts of southern Africa and from Paleozoic ultramafic complexes of the European Alps; metallogenic and geochemical implications. *Economic Geology* 77, 1–24.
- Santosh, M., 2010. Assembling North China Craton within the Columbia supercontinent: the role of double-sided subduction. *Precambrian Research* 178, 149–167.
- Santosh, M., Omori, S., 2008. CO₂ flushing: a plate tectonic perspective. *Gondwana Research* 13, 86–102.
- Santosh, M., Maruyama, S., Omori, S., 2009. A fluid factory in Solid Earth. *Lithosphere* 1, 29–33.
- Sarma, D.S., Fletcher, I.R., Rasmussen, B., McNaughton, N.J., Mohan, M.R., Groves, D.I., 2011. Archean gold mineralization synchronous with late cratonization of the Western Dharwar Craton, India: 2.52 Ga U–Pb ages of hydrothermal monazite and xenotime in gold deposits. *Mineralium Deposita* 46, 273–288.
- Seno, T., Kirby, S.H., 2014. Formation of plate boundaries: the role of mantle devolatilization. *Earth-Science Reviews* 129, 85–99.
- Sibson, R.H., 1992. Implications of fault-valve behaviour for rupture nucleation and recurrence. *Tectonophysics* 211, 282–293.
- Sibson, R.H., 2004. Controls on maximum fluid overpressure defining conditions for mesozonal mineralization. *Journal of Structural Geology* 26, 1127–1136.
- Sibson, R.H., 2013. Stress switching in subduction forearcs: implications for overpressure containment and strength cycling on megathrusts. *Tectonophysics* 600, 142–152.
- Song, M.-C., Li, S.-Z., Santosh, M., Zhao, S.-J., Yu, S., Yi, P.-H., Cui, S.-X., Lu, G.-X., Xu, J.-X., Song, Y.-X., Zhou, M.L., 2015. Types, characteristics and metallogenesis of gold deposits in the Jiaodong Peninsula, eastern North China craton. *Ore Geology Reviews* 65, 612–625.
- Standish, C.D., Dhuime, B., Chapman, R.J., Hawkesworth, C.J., Pike, A.W.G., 2014. The genesis of gold mineralization hosted by orogenic belts: a lead isotope investigation of Irish gold deposits. *Chemical Geology* 378–379, 40–51.
- Steadman, J.A., Large, R.R., Meffre, S., Bull, S.W., 2013. Age, origin, and significance of nodule sulfides in 2680 Ma carbonaceous black shale of the Eastern Goldfields Superterrane, Yilgarn craton, Western Australia. *Precambrian Research* 230, 227–247.
- Stuwe, K., 1998. Tectonic constraints on the timing relationships of metamorphism, fluid production and gold-bearing quartz vein emplacement. *Ore Geology Reviews* 13, 219–228.
- Thompson, J.F.H., Sillitoe, R.H., Baker, T., Lang, J.R., Mortensen, J.K., 1999. Intrusion-related gold deposits associated with tungsten-tin provinces. *Mineralium Deposita* 34, 323–334.
- Tomkins, A.G., 2010. Windows of metamorphic sulfur liberation in the crust: implications for gold deposit genesis. *Geochimica et Cosmochimica Acta* 74, 3246–3259.
- Tomkins, A.G., Grundy, C., 2009. Upper temperature limits of orogenic gold deposit formation: Constraints from the granulite-hosted Griffin's Find deposit, Yilgarn Craton. *Economic Geology* 104, 669–685.
- Tomkins, A.G., Mavrogenes, J.A., 2002. Mobilization of gold as a polymetallic melt during pelite anatexis at the Challenger deposit, South Australia—a metamorphosed Archean gold deposit. *Economic Geology* 97, 1249–1271.
- Tomkins, A.G., Pattison, D.R.M., Zaleski, E., 2004. The Hemlo gold deposit, Ontario: an example of melting and mobilization of a precious metal-sulfosalt assemblage during amphibolite facies metamorphism and deformation. *Economic Geology* 99, 1063–1084.
- Touret, J.L.R., Huizenga, J.M., 2012. Fluid-assisted granulite metamorphism: a continental journey. *Gondwana Research* 21, 224–235.
- Vielreicher, N.M., Groves, D.I., McNaughton, N.J., Fletcher, I.R., 2015. The timing of gold mineralization across the eastern Yilgarn using U–Pb geochronology of hydrothermal phosphate minerals. *Mineralium Deposita* 391–428.
- Vielreicher, N.M., Groves, D.I., Snee, L.W., Fletcher, I.R., McNaughton, N.J., 2010. Broad synchronicity of three gold mineralization styles in the Kalgoorlie gold field: SHRIMP, U–Pb, and ⁴⁰Ar/³⁹Ar geochronological evidence. *Economic Geology* 105, 187–227.
- Wall, V.J., Graupner, T., Yantsen, V., Seltmann, R., Hall, G.C., 2004. Muruntau, Uzbekistan—a giant thermal aureole gold (TAG) system. In: Muhling, J., Goldfarb, R., Vielreicher, N., Bierlein, F.P., Stumpff, E., Groves, D.I., Kenworthy, S. (Eds.), *SEG 2004: Predictive Mineral Discovery under Cover; Extended Abstracts*. Centre for Global Metallogeny, vol. 33. University of Western Australia, Publication, pp. 199–203.
- Walshe, J.L., Halley, S.W., Hall, G.A., Kitto, P., 2003. Contrasting fluid systems, chemical gradients and controls on large-tonnage, high-grade Au deposits, Eastern Goldfields Province, Yilgarn Craton, Western Australia. In: *Mineral Exploration and Sustainable Development*. 7th Biennial SGA meeting, Athens, pp. 827–830.
- Wang, L.-G., Qiu, Y.-M., McNaughton, N.J., Groves, D.I., Luo, Z.-K., Huang, J.-Z., Miao, L.-C., Liu, Y.-K., 1998. Constraints on crustal evolution and gold metallogeny in the northwestern Jiaodong peninsula, China, from SHRIMP U–Pb zircon studies of granitoids. *Ore Geology Reviews* 13, 275–291.
- Webber, A.P., Roberts, S., Taylor, R.N., Pitcairn, I.K., 2013. Golden plumes: substantial gold enrichment of oceanic crust during ridge-plume interaction. *Geology* 41, 87–90.
- White, A.J.R., Waters, D.J., Robb, L.J., 2015. Exhumation-driven devolatilization as a fluid source for orogenic gold mineralization at the Damang deposit, Ghana. *Economic Geology* 110, 1009–1026.
- Wilson, C.J.L., Schaubs, P.M., Leader, L.D., 2013. Mineral precipitation in the quartz reefs of the Bendigo gold deposit, Victoria, Australia. *Economic Geology* 108, 259–278.
- Windley, B.F., Maruyama, S., Xiao, W.-J., 2010. Delamination/thinning of subcontinental lithospheric mantle under eastern China: the role of water and multiple subduction. *American Journal of Science* 310, 1250–1293.
- Wyman, D., Kerrich, R., 2010. Mantle plume-volcanic arc interaction: consequences for magmatism, metallogeny, and cratonization in the Abitibi and Wawa subprovinces, Canada. *Canadian Journal of Earth Sciences* 47, 565–589.
- Wyman, D.A., O'Neill, C.O., Ayer, J.A., 2008. Evidence for Modern-style Subduction to 3.1Ga: a Plateau-adakite-gold (Diamond) Association, vol. 440. Geological Society of America Special Publication, pp. 129–148.
- Yamaguchi, A., Cox, S.F., Kimura, G., Okamoto, S., 2011. Dynamic changes in fluid redox state associated with episodic fault rupture along a megasplay fault in a subduction zone. *Earth and Planetary Science Letters* 302, 369–377.
- Yang, Q.Y., Santosh, M., 2015. Early Cretaceous magma flare-up and its implications on gold mineralization in the Jiaodong Peninsula, China. *Ore Geology Reviews* 65, 626–642.
- Yang, Q.Y., Santosh, M., Shen, J.F., Li, S.R., 2014. Juvenile vs. recycled crust in NE China: zircon U–Pb geochronology, Hf isotope and an integrated model for Mesozoic gold mineralization in the Jiaodong Peninsula. *Gondwana Research* 25, 1445–1468.
- Yang, L.-Q., Deng, J., Guo, R.-P., Guo, L.-N., Wang, Z.-L., Chen, B.-H., Wang, X.-D., 2016a. World-class Xincheng gold deposit: an example from the giant Jiaodong gold province. *Geoscience Frontiers* 7 (3), 419–430.
- Yang, L.-Q., Deng, J., Wang, Z.-L., Guo, L.-N., Li, R.-H., Groves, D.I., Danyushevskiy, J.V., Zhang, C., Zhang, X.-L., Zhao, H., 2016b. Relationships between gold and pyrite at the Xincheng gold deposit, Jiaodong Peninsula, China: implications for gold source and deposition in a brittle epizonal environment. *Economic Geology* 111 (1), 105–126.